

WELD Experiences



David P. Roy, Hankui K. Zhang, Petr Votava*, Adam Dosch, Jian Li

Geospatial Sciences Center of Excellence, Wecota Hall, South Dakota State University,
Brookings, SD 57007, USA

* NASA Ames Research Center, Moffett Field, CA 94035, USA

Landsat Analysis Ready Data Session

Winter Landsat Science Team Meeting,
Newman Library Multipurpose Room,
Virginia Tech, Blacksburg, VA
January 12-14 2016





Contents lists available at ScienceDirect

Remote Sensing of Environment

journal homepage: www.elsevier.com/locate/rse

Landsat-8: Science and product vision for terrestrial global change research



- The provision of ‘higher-level’ Landsat products, i.e., geographically seamless, gridded products that have been subject to geophysical transformations and processed to derive environmental variables over different time periods (monthly, seasonal, annual), have been advocated by the LST and by the user community.
- Higher-level products are needed to meet demands for consistently processed, moderate spatial resolution, large area, long-term terrestrial data records for climate and global change studies, to help national and international reporting linked to multilateral environmental agreements, and for regional and national resource management applications.

What the Landsat user community wants ...

Summary Slide from a David Roy presentation at the
Landsat Science Team Meeting, January 19-21, 2010

Computer History Museum, 1401 N Shoreline Boulevard, Mountain View, CA 94043

- Derived Landsat data products for free
- Systematic, consistent, community endorsed data processing
 - calibration, geolocation
 - radiometric normalization / BRDF correction, atmospheric correction
 - cloud-screened, snow-screened, SLC-off gap filling
 - needed in order to derive higher level bio/geophysical products
- Composited large-area data product mosaics
 - updated at the pixel level
 - using all the Landsat data, not just select acquisitions
 - processed shortly after acquisition i.e. “near real time”
- A long term Landsat data product record
- *Similar to the NASA MODIS land products but at high spatial resolution*
- *above is what the NASA WELD funded project is seeking to achieve*

WELD Tiles

5000 x 5000 30m pixels


Albers Coordinates


Landsat

6300 x 6300 30m pixels

UTM Coordinates

~3,100 million 30m land pixels






WELD Weekly Products

Week 30: July 22 - 28 2008



WELD Monthly Products

July 2008



Albers projection

WELD Seasonal Products

Summer (June, July, August) 2008



Albers projection

WELD Annual Products

(December 2007 - November 2008)



~ 10,000 L1T images / year


Albers projection

Available Years:

CONUS



Alaska




<http://weld.cr.usgs.gov>

CONUS 2008

<< Home


Annual & Seasonal



Monthly



Weekly



Version 1.5 WELD Product Format (all pixels are 30m)

MODIS like format and storage approach

Science Data Set Name	Data Type	Valid Range	Scale factor	Units	Fill Value
Band1_TOA_REF	int16	-32767 -- 32767	10000	Unitless	-32768
Band2_TOA_REF	int16	-32767 -- 32767	10000	Unitless	-32768
Band3_TOA_REF	int16	-32767 -- 32767	10000	Unitless	-32768
Band4_TOA_REF	int16	-32767 -- 32767	10000	Unitless	-32768
Band5_TOA_REF	int16	-32767 -- 32767	10000	Unitless	-32768
Band61_TOA_BT	int16	-32767 -- 32767	100	Degrees Celsius	-32768
Band62_TOA_BT	int16	-32767 -- 32767	100	Degrees Celsius	-32768
Band7_TOA_REF	int16	-32767 -- 32767	10000	Unitless	-32768
NDVI_TOA	int16	-10000 -- 10000	10000	Unitless	-32768
Day_Of_Year	int16	1 -- 366	1	Day	0
Saturation_Flag	uint8	0 -- 255	1	Unitless	None
DT_Cloud_State	uint8	0, 1, 2, 200	1	Unitless	255
ACCA_State	uint8	0, 1	1	Unitless	255
Num_Of_Observations	uint8	0 -- 255	1	Unitless	None

- Format designed to be easy to use for science and applications



- ☒ Vector
- Zoom In (or double click)
- Zoom Out (or mouse wheel)
- Next Period
- Previous Period

Order Coordinates

- ☒ Longitude/latitude
- ☐ Albers

Hold the shift button & drag the mouse to define your order area, or enter the area coordinates below

Note: In using Long/Lat the coordinates refer to the SW and NE corners of the area

North:

South:

West:


East:

Click the left button to define a single pixel time series dump, or enter the pixel coordinates below

North:

West:

WELD derived 30m Vegetation Continuous field product



Percent cover

Bare ground


Grass/shrubs/moss

Trees

Proof of Concept generated from
2008 monthly, seasonal and
annual WELD composites

Matt Hansen et al. RSL 2011

WELD derived 30m 5-year **forest cover loss** and **bare ground gain** product: 2006-2010



Landsat-MODIS fusion

Jan

Dec



Boschetti et al. 2015, RSE


Landsat-MODIS fusion

Jan

Dec




MTBS polygons



Boschetti et al. 2015, RSE


2010 CONUS crop field size map

(mean field size shown in 7.5 x 7.5 km grid cells)



derived from all 13,666 WELD processed Landsat 5 and 7 scenes
available in the U.S. Landsat archive for 12 months Yan and Roy, 2016, *RSE*

Global WELD code run on NASA Earth Exchange (NEX)



- 2 PB on-line storage, 50PB tape storage
- 512 cores dedicated cores
- 200,000 total available cores on Pleiades



Pleiades
NASA's fastest supercomputer



COLLABORATION
(over 400 members)

COMPUTING
(180,000 cores)

Data Repository
(over 600TB of data)

3 Years global 30m monthly & annual WELD Version 2.2 products available



30m Global WELD NEX

Month 12 2008

GeoTiff format products: <http://globalweld.cr.usgs.gov>

HDF format products: <http://globalweld.cr.usgs.gov/collections>

Climate year 2011


<< Home



Climate year 2010



Climate year 2009



Notice: To obtain large amounts of WELD data, please use the [DAAC2Disk Download Manager](#) to submit your request. This will ensure your order will process through our system efficiently. Please see the [DAAC2Disk User Guide](#) for more information and direct any questions to [LP DAAC User Services](#).

Collections

74TB

312,000 files


* The text files were created to be used with the DAAC2Disk tool. This tool facilitates the downloading of large amounts of data.

Annual (2009, 2010, 2011)				
Folder Name	Type	# of files	Size	Text File
weld.global.annual.2009	Global Annual 2009	8,050	3,026,904 MB	weld.global.annual.2009.txt
weld.global.annual.2010	Global Annual 2010	8,023	2,975,832 MB	weld.global.annual.2010.txt
weld.global.annual.2011	Global Annual 2011	8,003	2,943,370 MB	weld.global.annual.2011.txt
	Totals for Annuals	24,076	8,946,106 MB	

2009				
Folder Name	Type	# of files	Size	Text File
weld.global.month01.2009	Global Month 1	5,770	1,451,543 MB	weld.global.month01.2009.txt
weld.global.month02.2009	Global Month 2	6,197	1,528,338 MB	weld.global.month02.2009.txt
weld.global.month03.2009	Global Month 3	7,153	1,835,942 MB	weld.global.month03.2009.txt
weld.global.month04.2009	Global Month 4	7,233	1,777,753 MB	weld.global.month04.2009.txt
weld.global.month05.2009	Global Month 5	7,245	1,854,749 MB	weld.global.month05.2009.txt
weld.global.month06.2009	Global Month 6	7,104	1,953,828 MB	weld.global.month06.2009.txt
weld.global.month07.2009	Global Month 7	6,884	1,896,111 MB	weld.global.month07.2009.txt
weld.global.month08.2009	Global Month 8	7,018	1,940,278 MB	weld.global.month08.2009.txt
weld.global.month09.2009	Global Month 9	7,178	1,990,522 MB	weld.global.month09.2009.txt
weld.global.month10.2009	Global Month 10	7,095	1,858,060 MB	weld.global.month10.2009.txt
weld.global.month11.2009	Global Month 11	6,278	1,657,394 MB	weld.global.month11.2009.txt
weld.global.month12.2008	Global Month 12	5,511	1,341,593 MB	weld.global.month12.2008.txt
	Totals for Months	80,666	21,086,111 MB	


2010				
Folder Name	Type	# of files	Size	Text File
weld.global.month01.2010	Global Month 1	5,605	1,396,719 MB	weld.global.month01.2010.txt
weld.global.month02.2010	Global Month 2	6,117	1,445,984 MB	weld.global.month02.2010.txt
weld.global.month03.2010	Global Month 3	6,082	1,506,701 MB	weld.global.month03.2010.txt

Global Version 2.2 WELD derived 30m Vegetation Continuous field product




(Matt Hansen, Peter Potapov et al.)


Overview of Global **Version 2.2** WELD Processing Sequence



Overview of Global **Version 3.0** WELD Processing Sequence



Overview of Global Version 3.0 WELD Processing Sequence









UAT




Active



BASE LAYERS

-  Corrected Reflectance (True Color, Monthly, v3, NEX, Standard) 
WELD_CR321_Mon_LL_STD - Landsat
-  NDVI (Monthly, v3, NEX, Standard) 
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



2000 km
2000 mi

2009 JUN 01



DAYS

MONTHS

YEARS





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



1000 m

5000 ft

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



2 km

2 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS

VIEW





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



5 km

5 mi

2009 JUN 01



MAY 2009



JUN 2009

JUL 2009

DAYS

MONTHS





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



10 km

5 mi



2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat



OVERLAYS



20 km

10 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



50 km

20 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS

YEARS





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



+

-

100 km

50 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS

YEARS





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



200 km

100 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS

YEARS



UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



500 km

200 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS

YEARS



UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



500 km
500 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS





UAT




Active



BASE LAYERS

	Corrected Reflectance (True Color, Monthly, v3, NEX, Standard) WELD_CR321_Mon_LL_STD - Landsat	
	NDVI (Monthly, v3, NEX, Standard) WELD_NDVI_Mon_LL_STD - Landsat	

OVERLAYS



1000 km
1000 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS

WEEKS





UAT




Active



BASE LAYERS

	Corrected Reflectance (True Color, Monthly, v3, NEX, Standard) WELD_CR321_Mon_LL_STD - Landsat	
	NDVI (Monthly, v3, NEX, Standard) WELD_NDVI_Mon_LL_STD - Landsat	

OVERLAYS



2000 km

2000 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS

WEEK





UAT




Active



BASE LAYERS

	Corrected Reflectance (True Color, Monthly, v3, NEX, Standard) <i>WELD_CR321_Mon_LL_STD - Landsat</i>	
	NDVI (Monthly, v3, NEX, Standard) <i>WELD_NDVI_Mon_LL_STD - Landsat</i>	

OVERLAYS



5000 km
2000 mi

2009 JUN 01




DAYS

MONTHS


WEEK




Overview of Global Version 3.0 WELD Processing Sequence



Overview of Global Version 3.0 WELD Processing Sequence



WELD Landsat 5 & 7 TOA reflectance



Global WELD NEX Version 3.0 Annual 2009 30m product
from 141,098 L1T scenes (59,183 Landsat 5 & 81,915 Landsat 7)

Sinusoidal Equal Area Projection


WELD Landsat 5 & 7 surface reflectance normalized BRDF adjusted reflectance composite



Global WELD NEX Version 3.0 Annual 2009 30m product
from 141,098 L1T scenes (59,183 Landsat 5 & 81,915 Landsat 7)

Sinusoidal Equal Area Projection


WELD Landsat 5 & 7 TOA NDVI NBAR



Global WELD NEX Version 3.0 Annual 2009 30m product
from 141,098 L1T scenes (59,183 Landsat 5 & 81,915 Landsat 7)

Sinusoidal Equal Area Projection

WELD Landsat 5 & 7 surface NDVI NBAR



Global WELD NEX Version 3.0 Annual 2009 30m product
from 141,098 L1T scenes (59,183 Landsat 5 & 81,915 Landsat 7)

Sinusoidal Equal Area Projection



UAT




Active



BASE LAYERS

	Corrected Reflectance (True Color, Monthly, v3, NEX, Standard) WELD_CR321_Mon_LL_STD - Landsat	
	NDVI (Monthly, v3, NEX, Standard) WELD_NDVI_Mon_LL_STD - Landsat	

OVERLAYS



Global WELD
June 2009
Surface NBAR

20 km
10 mi

2009 JUN 01



DAYS

MONTHS





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



Global WELD
June 2009
Surface NDVI

20 km

10 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS

YEARS



UAT



Active



BASE LAYERS

	Corrected Reflectance (True Color, Monthly, v3, NEX, Standard) WELD_CR321_Mon_LL_STD - Landsat	
	NDVI (Monthly, v3, NEX, Standard) WELD_NDVI_Mon_LL_STD - Landsat	

OVERLAYS



1000 m
5000 ft

2009 JUN 01



DAYS
MONTHS





UAT



Active



BASE LAYERS

	Corrected Reflectance (True Color, Monthly, v3, NEX, Standard) WELD_CR321_Mon_LL_STD - Landsat	
	NDVI (Monthly, v3, NEX, Standard) WELD_NDVI_Mon_LL_STD - Landsat	

OVERLAYS



2 km
2 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



5 km

5 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS

WIDE





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



10 km

5 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS





UAT



Active



BASE LAYERS

	Corrected Reflectance (True Color, Monthly, v3, NEX, Standard) WELD_CR321_Mon_LL_STD - Landsat	
	NDVI (Monthly, v3, NEX, Standard) WELD_NDVI_Mon_LL_STD - Landsat	

OVERLAYS



20 km
10 mi

2009 JUN 01



DAYS

MONTHS





UAT



Active



BASE LAYERS

	Corrected Reflectance (True Color, Monthly, v3, NEX, Standard) WELD_CR321_Mon_LL_STD - Landsat	
	NDVI (Monthly, v3, NEX, Standard) WELD_NDVI_Mon_LL_STD - Landsat	

OVERLAYS



50 km

20 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS





UAT



Active



BASE LAYERS

	Corrected Reflectance (True Color, Monthly, v3, NEX, Standard) WELD_CR321_Mon_LL_STD - Landsat	
	NDVI (Monthly, v3, NEX, Standard) WELD_NDVI_Mon_LL_STD - Landsat	

OVERLAYS



100 km
50 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



200 km

100 mi

2009 JUN 01



MAY 2009

JUN 2009

JUL 2009

DAYS

MONTHS

YEARS





UAT



Active



BASE LAYERS



Corrected Reflectance (True Color, Monthly, v3, NEX, Standard)
WELD_CR321_Mon_LL_STD - Landsat



NDVI (Monthly, v3, NEX, Standard)
WELD_NDVI_Mon_LL_STD - Landsat

OVERLAYS



500 km

200 mi

2009 JUN 01



MAY 2009

JUN 2009


JUL 2009

DAYS


MONTHS



Overview of Global Version 3.0 WELD Processing Sequence



WELD Landsat 5 & 7 surface reflectance one week composite




Arizona
500 x 500 30m pixels



Mississippi

WELD Landsat 5 & 7 surface reflectance one week composite

normalized BRDF adjusted reflectance (NBAR)
0° view zenith & local overpass solar zenith




Arizona
500 x 500 30m pixels



Mississippi

c-factors over MODIS 110° FOV


derived using mean MDC43 spectral BRDF model parameters



Roy et al. A general method to normalize Landsat reflectance data to nadir BRDF adjusted reflectance, *Remote Sensing of Environment*.

c-factors over Landsat 15° FOV

derived using mean MDC43 spectral BRDF model parameters




Roy et al. A general method to normalize Landsat reflectance data to nadir BRDF adjusted reflectance, *Remote Sensing of Environment*.

c-factors over MODIS 110° FOV

derived using mean MDC43 spectral BRDF model parameters


(similar over Sentinel-2 20.6° FOV)



Roy et al. A general method to normalize Landsat reflectance data to nadir BRDF adjusted reflectance, *Remote Sensing of Environment*.

WELD Landsat 5 & 7 surface reflectance one week composite

normalized BRDF adjusted reflectance (NBAR)
0° view zenith & local Landsat overpass solar zenith




Arizona
500 x 500 30m pixels



Mississippi

12 months of non-Antarctic (147,358 Landsat 5 and 7 acquisitions)
Landsat solar zenith (θ_s) plotted as a function of scene center latitude and day of year




5th degree polynomial used to model the local overpass time as a function of latitude for both Landsat sensors:

$$\hat{t}_{local} = 1.36292 \times 10^{-9} \alpha^5 - 3.15403 \times 10^{-8} \alpha^4 - 3.15819614 \times 10^{-6} \alpha^3 + 0.0000652685643 \alpha^2 + 0.0120604786763 \alpha + 10.06$$

Zhang, H.K., Roy, D.P., & Kovalskyy, V. (2016). Optimal solar geometry definition for global long-term Landsat time-series bidirectional reflectance normalization. *IEEE TGRS*, *In press*.


WELD Landsat 5 & 7 observed Solar Zenith



Global WELD NEX Version 3.0 September 2009 30m product
from 15,058 L1T scenes (7,328 Landsat 5 & 7,730 Landsat 7)

Sinusoidal Equal Area Projection

WELD Landsat 5 & 7 modelled Solar Zenith used to derived the WELD NBAR




Global WELD NEX Version 3.0 September 2009 30m product
from 15,058 L1T scenes (7,328 Landsat 5 & 7,730 Landsat 7)

Sinusoidal Equal Area Projection

Long term Landsat NBAR generation implications ...

Landsat 5 local time derived from Landsat metadata



Landsat
path/row 27/42
(somewhere in Texas)

Local time difference
0.9714 hours over 27
years of Landsat 5 !

WELD Landsat 5 & 7 surface reflectance normalized BRDF adjusted reflectance composite




Global WELD NEX Version 3.0 Annual 2009 30m product
from 141,098 L1T scenes (59,183 Landsat 5 & 81,915 Landsat 7)

Sinusoidal Equal Area Projection

Global WELD
Version 3.0
(NBAR
surface
reflectance)
available
February
2016


Planned Global WELD V3.0 Production

Monthly + Annual 30m products, fusion of contemporaneous Landsat 4,5,7



Landsat repatriation from other space agencies will provide more Landsat data in earlier epochs

USGS archive prior to LGAC




Number of images



Archive holdings pre-LGAC

Wulder, et al. 2016. RSE.

USGS archive up to Jan 1, 2015




Number of images



Archive holdings as of
January 1, 2015

Wulder, et al. 2016. RSE.

Future archive once outstanding images added



Number of images



Potential future archive holdings (with LGAC)

Wulder, et al. 2016. RSE.

Some Lessons Learned From Bulk Landsat Science Data Processing

Summary Slide from a David Roy presentation at the

Landsat Science Team, USGS EROS Center, Sioux Falls, August 16-18 2011

- You cannot hit a *big red button* to make reliable large area and/or long term Landsat higher level science data products
- Get the product out early, Ensure mechanisms to involve the user community in product QA
- Routinely QA the input data with validation
- Reprocess products as needed
- Implement and manage a product versioning scheme
- Science algorithms developed on individual scenes don't necessarily provide consistency between scenes
- Product Quality Assessment challenge increases with data volume, algorithm complexity and product dependencies
- Expect schedule creep in Science Product Development

Version 3 NEX WELD rejections

- PERIOD : 07_2009
- ALL L5+L7:
- NUMBER STAGED SCENES : 20309
- NUMBER REJECTED SCENES (NOT L1T) : 4763
- NUMBER REJECTED (RMSE > 30m) : 49
- NUMBER REJECTED (SOLAR ZENITH > 85): 0
- NUMBER SCENES CONSIDERED/PROCESSED : 15497



All is well ! A relief that the _GEOM file was not produced for this mutant (see attached) its Eastern edge is not parallel to its Western edge and it has a non-standard size (7091 x 7311 pixels). No way sh/could the code handle this.

If the _GEOM file does not get produced then the UTM image will not get used in the Tile processing.

Petr, if its easy would be interesting to know what percentage of a global month has “The R-sq of edge fitting is only ...” errors.

Hank, tomorrow please check L457.geometry.v1.5.c and see what the R2 threshold is to exit the process (this was something Junchang and I worked on pre-Val).

Science needs for derived Higher Level products

- Algorithms

- recent, peer-reviewed, responsive to science community requirements
- taking advantage of improved sensing technology

- Standardized products

- format
- filename convention
- content structure (e.g., data layers, metadata etc.)
- documentation
- versioning (perhaps also collections)

- Dedicated scientists & support staff to

- develop, code, and document algorithms & products
- monitor/define product quality and product accuracy
- refine algorithms as needed
- develop software tools for product manipulation
- weekly telecons

- Systematic production on dedicated computing facility, allowing

- long term global production
- sophisticated pre-processing and use of ancillary data sets
- reprocessing to achieve stability of products over long time periods
- production efficiencies (e.g., for products that share common inputs)

Climate year 2011

<< Home

Annual 2011

Climate year 2010

Annual 2010

Climate year 2009

Annual 2009

December 2008

January 2009


February 2009

March 2009

April 2009

May 2009

June 2009



July 2009

August 2009

September 2009

October 2009

November 2009

Global
WELD
Version 3.0
(NBAR
surface
reflectance)
available
February
2016

Notice: These Version 3.0 products have replaced the Version 2.2 products. Version 3.0 is only available for viewing and on-demand ordering on this WYSIWYG interface, however, both versions are available as [HDF tiled products](#) for download.

Notice: To obtain large amounts of WELD data, please use the [DAAC2Disk Download Manager](#) to submit your request. This will ensure your order will process through our system efficiently. Please see the [DAAC2Disk User Guide](#) for more information and direct any questions to [LP DAAC User Services](#).

Landsat 8
red, green, blue
TOA reflectance


Cape Town,
South Africa

WELD week 49
(December 3 - 9 2015)

2 input L1T images
sensed:
December 8
08:35:04
08:35:28

nearest neighbor
resampled to
global WELD tile
hh19vv12h3v2

5295 x 5295 30m pixels



Sentinel 2A
red, green, blue
TOA reflectance

Cape Town,
South Africa

WELD week 49
(December 3 - 9 2015)

1 input L1C image
sensed:
December 8
08:50:36

box car
resampled to
global WELD tile
hh19vv12h3v2

5295 x 5295 30m pixels

